

Name: _____ Section: _____ Date: _____

Worksheet - Exp 22: Thin Lenses

Objective

The objective of this experiment is to measure the focal lengths of a converging lens and a diverging lens and investigate magnification.

Theory

Light refracts (bends) when passing through media with different indices of refraction. This property can be very useful, especially when a *thin lens* is used. A thin lens' thickness is much less than its diameter.

A converging (convex, positive) lens is thicker in the center than at the edges. It can be used to focus parallel light rays and form a *real image*. A real image can be projected on a screen. The image exists regardless of whether or not a screen is in position to show it.

A diverging (concave, negative) lens normally forms a *virtual image*. Light rays do not actually pass through a virtual image. It cannot be projected on a screen. When you look at yourself in a mirror, you are looking at a virtual image. If the object is real, the image is virtual (with a diverging lens). However, when a diverging lens is used in combination with a converging lens, for instance, the object can be virtual, the image real. Parameters must be met for a real image to be formed; read the *Part 2* procedure carefully.

An important property of a lens is its focal length, f . The focal length of a thin lens is given by:

$$\frac{1}{d_o} + \frac{1}{d_i} = \frac{1}{f}$$

where d_o is the object distance and d_i is the image distance. These distances are measured **from the lens**.

PROCEDURE


Part 1: Converging Lens

Method I - Use the lens equation ($\frac{1}{d_o} + \frac{1}{d_i} = \frac{1}{f}$)

1. Mount the lens, screen, and light source on the optical bench. Adjust the height of the object, lens(es), and screen so that the optical axis passes through the center of each element.
2. Adjust the position of the lens and the screen until a clear image of the object is projected onto the screen. Considering the lens equation, how many combinations of d_i and d_o are possible?
3. Using the diagram to the right, record the position of each device: O , L , i . Positions are measured directly from the optics bench; a line is scribed on each holder for accuracy.
4. Calculate d_o , d_i , and f .

$$d_o = L - O$$

$$d_i = i - L$$



(6 pts)

O is the position of the object.

L is the position of the lens.

i is the position of the image.

d_o is the calculated object distance (absolute value).

d_i is the calculated image distance (absolute value).

Method II - Use a distant object ($d_o \rightarrow \infty$)

5. Hold lenses carefully by the edge. Project the image of a distant object on a screen. One way to achieve this is to take the lens and a ruler to a long hallway. Hold the lens such that light from a distant light source at the other end of the hallway passes through the lens and focuses on the wall.
6. Adjust the distance between the lens and the screen until a clear, distinct image of the distant light source is projected onto the screen.
7. Measure the distance from the lens to the screen. This distance is f . The lens equation shows that when d_o is large, $d_i \rightarrow f$.


$$f = \underline{\hspace{2cm}}$$

8. Is the image inverted? Magnified? Reversed? (4 pts)
9. Compare f (% difference) from the two methods. (4 pts)
10. Draw a ray diagram for Method I. Use at least two rays. (8 pts)
11. Draw a ray diagram for Method II. Use at least two rays. (8 pts)

Part 2: Diverging Lens

To determine the focal length of a diverging lens, the lens must create a measurable, real image as in *Part 1*. However, light cannot be focused through a diverging lens to form a real image unless that light was already converging. To accomplish this, a *real image* from a converging lens will be used as a *virtual object*.²

12. Form a real image using a converging lens. *Note:* d_o should be greater than $2f$.
13. Place the diverging lens between the converging lens and its (real) image; refer to the figure below. The *real image* from the converging lens is now a *virtual object* for the diverging lens.
14. Determine the position of the diverging lens' image by adjusting the screen's position.



15. Record your data in the digram above.
16. Determine d_o and d_i for the diverging lens.
17. Calculate f for the diverging lens.

Converging Lens:
 O: _____ L: _____ i: _____
 d_o : _____ d_i : _____

Diverging Lens:
 L: _____ O: _____ i: _____
 d_o : _____ d_i : _____
 f_1 = _____ f_2 = _____

(12 pts)

Compare f of a Diverging Lens

18. Recalculate f using the following equation:

$$f = \frac{VW}{V-W}$$

where V and W are defined as:

$$V \equiv |d_o| \quad \text{and} \quad W \equiv |d_i|$$

(6 pts)

19. Compare f values from the lens equation and the equation above for the diverging lens. If you followed the sign conventions closely, the f values should be identical.

²For a diverging lens, either the object or the image can be real; the other must be virtual.

20. Consider a *concave* lens made out of air that is immersed in water (perhaps two watch glasses glued to each end of a piece of pipe, with air inside). Will it form a real image that can be focused on a screen? Draw a ray diagram to support your answer. (8 pts)

Part 3: Lateral Magnification, M

Use only the converging lens to investigate lateral magnification. Record all data in the diagrams to the right.

21. Set $d_o > 2f$ by adjusting the distance between the object and the lens. Find the image using your large screen. Record the position of the image and mark on the screen the top and bottom of the image.

22. Measure the image height, h_i , and the object height, h_o . Be sure to measure the same dimension on both object and image. (2 pts)

h_i : _____ h_o : _____

23. Calculate M :

$$M = \frac{h_i}{h_o} \quad M = \text{_____} \quad (2 \text{ pts})$$

24. Verify the magnification (M) using the following equation:


$$M = \frac{-d_i}{d_o} \quad M = \text{_____} \quad (2 \text{ pts})$$

25. Compare the two values of M .

% diff: _____ (2 pts)

Converging Lens: $d_o > 2f$

O: _____ L: _____ i: _____



d_o : _____ d_i : _____


f = _____

(6 pts)

26. Set $f < d_o < 2f$. Locate the image.
27. Measure h_i and h_o . (2 pts)
 h_i : _____ h_o : _____
28. Calculate M using both of the previous equations and compare the two values. (4 pts)
 $M_1 =$ _____ $M_2 =$ _____
 % diff: _____
29. Set $d_o = f$. Try to find d_i . Consider Method II and Eq. 21.1; where should the image be? (4 pts)
30. Set $d_o < f$. Look *through* the lens at the object. Note your observations. (4 pts)

Converging Lens: $f < d_o < 2f$

O: _____ L: _____ i: _____



d_o : _____ d_i : _____

$f =$ _____

(6 pts)

31. Set the lens carefully aside where it will not be damaged. Unplug the light source and lay the power cord across the optics bench.

32. If a convex lens with $n = 1.30$ and $f = 25$ cm is immersed in a fluid with an index of refraction that is also 1.30, what is the new focal length of the lens? Draw a ray diagram. (10 pts)