

PHYS 621 – LEGENDRE FUNCTIONS & SPHERICAL HARMONICS

LEGENDRE FUNCTIONS

Differential equation

$$(1 - z^2)u''(z) - 2zu'(z) + \left(\nu(\nu + 1) - \frac{\mu^2}{1 - z^2} \right)u(z) = 0,$$

Solution in the region $|z| < 1$:

$$P_\nu^\mu(z) = \frac{1}{\Gamma(1 - \mu)} \left(\frac{z + 1}{z - 1} \right)^{\mu/2} F(-\nu, \nu + 1; 1 + \mu; (1 - z)/2),$$

is called Legendre function of the first kind. It is also represented as

$$P_\nu^\mu(z) = \frac{2^\mu}{\Gamma(1 - \mu)} (z^2 - 1)^{-\mu/2} F(1 - \mu + \nu, -\mu - \nu; 1 - \mu; (1 - z)/2).$$

Note the property:

$$P_\nu^\mu(z) = P_{-\nu-1}^\mu(z).$$

For $\mu = 0$:

$$P_\nu(z) \equiv P_\nu^0(z) = F(-\nu, \nu + 1; 1; (1 - z)/2).$$

are the Legendre functions, which reduce to the Legendre polynomials for $\nu = l$ (integer). They satisfy the differential equation

$$(1 - z^2)P_\nu''(z) - 2zP_\nu'(z) + \nu(\nu + 1)P_\nu(z) = 0,$$

If $\mu \neq m$ (integer), $P_\nu^\mu(z)$ and $P_\nu^{-\mu}(z)$ are independent. Instead of $P_\nu^{-\mu}(z)$ it is customary to introduce:

$$Q_\nu^\mu(z) = \frac{\pi e^{i\pi\mu}}{2 \sin \pi\mu} \left(P_\nu^\mu(z) - \frac{\Gamma(\nu + \mu + 1)}{\Gamma(\nu - \mu + 1)} P_\nu^{-\mu}(z) \right).$$

(Legendre function of the second kind.)

If $\mu = m$ (integer) $P_\nu^m(z)$ and $P_\nu^{-m}(z)$ are proportional:

$$P_\nu^{-m}(z) = (-1)^m \frac{\Gamma(\nu - m + 1)}{\Gamma(\nu + m + 1)} P_\nu^m(z).$$

and

$$P_\nu^{-m}(z) = \frac{1}{\Gamma(1 + m)} \left(\frac{z + 1}{z - 1} \right)^{-m/2} F(-\nu, \nu + 1; 1 + m; (1 - z)/2),$$

Relation between $P_\nu^m(z)$ and the Legendre functions $P_\nu(z)$:

$$P_\nu^m(z) = (z^2 - 1)^{m/2} \frac{d^m}{dz^m} P_\nu(z), \quad m > 0.$$

ASSOCIATED LEGENDRE FUNCTIONS

If $\nu = l$ (integer) the associated Legendre functions are regular in $-1 \leq x \leq 1$. They are defined as:

$$P_l^m(x) = (-1)^m (1-x^2)^{m/2} \frac{d^m}{dx^m} P_l(x), \quad m > 0$$

Note the extra $(-1)^{3m/2}$ factor (conventions). Sometimes a normalization without $(-1)^m$ is also used.

Rodrigues' formula for the associated Legendre Polynomials:

$$P_l^m(x) = \frac{(-1)^m}{2^l l!} (1-x^2)^{m/2} \frac{d^{m+l}}{dx^{m+l}} (x^2-1)^l, \quad m > 0$$

Relation to $m < 0$:

$$P_l^{-m}(x) = (-1)^m \frac{(l-m)!}{(l+m)!} P_l^m(x).$$

Normalization:

$$\int_{-1}^1 dx P_l^m(x) P_{l'}^m(x) = \frac{2}{2l+1} \frac{(l+m)!}{(l-m)!} \delta_{ll'}.$$

This implies:

$$|m| < l, \quad \text{or} \quad -l \leq m \leq l.$$

SPHERICAL HARMONICS

Differential equation:

$$\left[\frac{\partial^2}{\partial \theta^2} + \cotan \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} + l(l+1) \right] Y(\theta, \varphi) = 0,$$

The regular solution in $-1 \leq \cos \theta \leq 1$ with boundary condition

$$Y(\theta, \varphi) = Y(\theta, \varphi + 2\pi)$$

is

$$Y_{lm}(\Omega) = \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} P_l^m(\cos \theta) e^{im\varphi}.$$

or

$$Y_{lm}(\Omega) = \frac{1}{2^l l!} \left[\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!} \right]^{1/2} (-\sin \theta)^m \frac{d^{m+l}}{d(\cos \theta)^{m+l}} (\cos^2 \theta - 1)^l e^{im\varphi}.$$

Normalization:

$$\int_{S^2} d\Omega Y_{lm}^* Y_{l'm'} = \delta_{ll'} \delta_{mm'}.$$

Symmetry property:

$$Y_{l,-m}(\Omega) = (-1)^m Y_{lm}^*(\Omega).$$

Completeness:

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l Y_{lm}^*(\Omega') Y_{lm}(\Omega) = \delta(\cos \theta - \cos \theta') \delta(\varphi - \varphi').$$

Addition theorem:

$$P_l(\cos \gamma) = \frac{4\pi}{2l+1} \sum_{m=-l}^l Y_{lm}^*(\Omega') Y_{lm}(\Omega),$$

where

$$\cos \gamma = \cos \theta' \cos \theta + \sin \theta' \sin \theta \cos(\varphi - \varphi').$$