

## Physics 651: Exercise 2

(not for submission)

1. The kets  $|u\rangle$ ,  $|v\rangle$ , and  $|w\rangle$  belong to a vector space that is spanned by the orthonormal basis  $\{|b_i\rangle\}$ . Let  $\hat{P} = \sum_{i,j} |b_i\rangle P_{i,j} \langle b_j|$  and  $\hat{Q} = \sum_{i,j} |b_i\rangle Q_{i,j} \langle b_j|$  be linear operators acting on that space. Which of the following expressions is incorrect?

(a)  $\langle u|\hat{P}|v\rangle^* = \langle v|\hat{P}^\dagger|u\rangle$

(b)  $(|u\rangle\langle v|)|w\rangle = \langle u|v\rangle|w\rangle$

(c)  $\langle b_i|\hat{P}\hat{Q}|u\rangle = \sum_{j,k} P_{i,j} Q_{j,k} u_k$

(d)  $(\hat{P}\hat{Q}|u\rangle)^\dagger = \langle u|\hat{Q}^\dagger\hat{P}^\dagger$

2. Let  $\alpha$ ,  $\beta$ , and  $\gamma$  be complex numbers and  $|u\rangle$ ,  $|v\rangle$ , and  $|w\rangle$  be elements of a complex vector space. Which of the following expressions is correct?

(a)  $(|u\rangle\langle v|)|w\rangle = \langle v|w\rangle|u\rangle$

(b)  $\langle u|(|v\rangle\langle w|) = \langle u|v\rangle^*|w\rangle$

(c)  $(\alpha|u\rangle \otimes |v\rangle \otimes |w\rangle)^\dagger = \alpha^* \langle w| \otimes \langle v| \otimes \langle u|$

(d)  $(\alpha|u\rangle + \beta|v\rangle + \gamma|w\rangle)^\dagger = \alpha\langle u| + \beta\langle v| + \gamma\langle w|$

3. Associated with a quantum system in its ground state  $|\psi\rangle$  is a density operator  $\hat{\rho} = |\psi\rangle\langle\psi|$ . When expressed in terms of a particular basis  $\{|n\rangle\}$ , the ground state has component amplitudes  $\psi_n = \langle n|\psi\rangle$ . For an observable  $\hat{O}$ , having matrix elements  $\langle m|\hat{O}|n\rangle = O_{m,n}$ , the ground state expectation value is

$$\langle \hat{O} \rangle = \frac{\text{tr } \hat{\rho} \hat{O}}{\text{tr } \hat{\rho}}.$$

Show that this is equivalent to

$$\frac{\sum_{m,n} \psi_m^* O_{m,n} \psi_n}{\sum_k |\psi_k|^2}.$$

4. The determinant of a  $2 \times 2$  matrix  $A$  is given by

$$\det A = \sum_{i=1}^2 \sum_{j=1}^2 \epsilon_{i,j} A_{1,i} A_{2,j}.$$

What is the correct definition of the alternating symbol?

(a)  $\epsilon_{1,1} = \epsilon_{2,2} = 0$  and  $\epsilon_{1,2} = \epsilon_{2,1} = 1$

(b)  $\epsilon_{1,1} = \epsilon_{2,2} = 0$  and  $\epsilon_{1,2} = -\epsilon_{2,1} = 1$

(c)  $\epsilon_{1,1} = \epsilon_{2,2} = 1$  and  $\epsilon_{1,2} = \epsilon_{2,1} = -1$

(d)  $\epsilon_{1,1} = -\epsilon_{2,2} = 1$  and  $\epsilon_{1,2} = \epsilon_{2,1} = 0$

5. The determinant of a  $4 \times 4$  matrix  $A$  is given by

$$\det A = \sum_{i=1}^4 \sum_{j=1}^4 \sum_{k=1}^4 \sum_{l=1}^4 \epsilon_{i,j,k,l} A_{1,i} A_{2,j} A_{3,k} A_{4,l},$$

where  $\epsilon_{i,j,k,l}$  is the 4-index Levi-Civita symbol. Which one of the following terms appears in the sum.

- (a)  $+A_{1,1}A_{2,2}A_{3,4}A_{4,3}$
- (b)  $-A_{1,3}A_{2,1}A_{3,4}A_{4,2}$
- (c)  $+A_{1,1}A_{2,2}A_{3,1}A_{4,2}$
- (d)  $-A_{1,3}A_{2,3}A_{3,3}A_{4,3}$

6. Here,  $|u\rangle$  and  $|v\rangle$  are elements of a vector space;  $\hat{A}$ ,  $\hat{B}$ , and  $\hat{C}$  are linear operators acting on the space; and  $\{|i\rangle\}$  constitutes an orthonormal basis for the space. Use the technique of inserting representations of unity,  $\hat{1} = \sum_i |i\rangle\langle i|$ , to prove that

$$\langle u | \hat{A} \hat{B} \hat{C} | v \rangle^* = \langle v | \hat{C}^\dagger \hat{B}^\dagger \hat{A}^\dagger | u \rangle.$$

7. Rotation about the  $x$ ,  $y$ , and  $z$  axes (in the right-hand sense about the directions  $\mathbf{e}_1$ ,  $\mathbf{e}_2$ , and  $\mathbf{e}_3$ ) is implemented by matrices

$$R_1(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{pmatrix}, R_2(\theta) = \begin{pmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{pmatrix}, R_3(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

- (a) Show that  $\det R_i(\theta) = 1$  for each of  $i = 1, 2, 3$  and for all values of the angle  $\theta$ .
- (b) Prove that  $R_i(-\theta) = R_i(\theta)^T = R_i(\theta)^{-1}$ .
- (c) Evaluate these three composite rotations:

$$\begin{aligned} A &= R_1(-\pi/2)R_2(\pi/2)R_1(\pi/2), \\ B &= R_3(\pi/2)R_2(\pi/4)R_1(\pi/2), \\ C &= R_1(-\pi/4)R_3(\pi/2)R_1(\theta)R_3(-\pi/2)R_1(\pi/4). \end{aligned}$$

In other words, evaluate each of the matrix products to determine the resulting  $3 \times 3$  matrix.

- (d) Prove that  $A$  corresponds to a rotation about  $\mathbf{e}_3$ ;  $B$  to a rotation about  $\mathbf{e}_1 + (1 + \sqrt{2})\mathbf{e}_2 + \mathbf{e}_3$ ; and  $C$  to a rotation about  $-\mathbf{e}_2 + \mathbf{e}_3$ . To determine the axes of rotation, solve the eigenproblems for  $(A + A^T)/2$ ,  $(B + B^T)/2$ , and  $(C + C^T)/2$ . Make use of  $\text{tr } A = 1 + 2 \cos \theta_A = 1$  and  $\text{tr } B = 1 + 2 \cos \theta_B = 1/\sqrt{2}$  to determine the angles of rotation. Evaluate  $\text{tr } C = 1 + 2 \cos \theta$  to confirm that the parameter  $\theta$  does in fact represent the rotation angle.

---

```

R1[\[Theta]_] := { { 1, 0, 0}, {0, Cos[\[Theta]], -Sin[\[Theta]]}, {0, Sin[\[Theta]],
  Cos[\[Theta]]} }
R2[\[Theta]_] := { { Cos[\[Theta]], 0, Sin[\[Theta]]}, {0, 1, 0}, {-Sin[\[Theta]], 0,
  Cos[\[Theta]]} }
R3[\[Theta]_] := { { Cos[\[Theta]], -Sin[\[Theta]], 0}, {Sin[\[Theta]], Cos[\[Theta]],
  0}, {0, 0, 1} }
A = R1[-\[Pi]/2] . R2[\[Pi]/2] . R1[\[Pi]/2]
B = R3[\[Pi]/2] . R2[\[Pi]/4] . R1[\[Pi]/2]
CC = R1[-\[Pi]/4] . R3[\[Pi]/2] . R1[\[Theta]] . R3[-\[Pi]/2] . R1[\[Pi]/4]
Solve[Tr[A] == 1 + 2 Cos[\[Theta]], \[Theta]]
Solve[Tr[A] == 1 + 2 Cos[\[Theta]], \[Theta]]
Solve[Tr[A] == 1 + 2 Cos[\[Theta]], \[Theta]]
EVA = Simplify[Eigensystem[(A + Transpose[A])/2]]
EVB = Simplify[Eigensystem[(B + Transpose[B])/2]]

```

```

EVC = Simplify[Eigensystem[(CC + Transpose[CC])/2]]
MemberQ[EVA[[1]], 1]
For[i = 1, i <= 3, ++i, If[EVA[[1]][[i]] == 1, Print[EVA[[2]][[i]]]]]
MemberQ[EVB[[1]], 1]
For[i = 1, i <= 3, ++i, If[EVB[[1]][[i]] == 1, Print[EVB[[2]][[i]]]]]
MemberQ[EVC[[1]], 1]
For[i = 1, i <= 3, ++i, If[EVC[[1]][[i]] == 1, Print[EVC[[2]][[i]]]]]

```

---

(e) Reflection across the  $x = 0$  plane is represented by the matrix

$$M = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

This maps every column vector

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} \text{ to a reflected vector } \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -x \\ y \\ z \end{pmatrix}.$$

Use a similarity transformation to determine the matrix  $M' = U^{-1}MU$  that reflects across the plane defined by  $y = -x$ .